FORM PTO-1449 (Modified) Department of Commerce tent and Trademark Office

Attorney Docket No.: STRATA-06949

Serial No.: 10/087,346

INFORMATION DISCLOSURE SHITEMENT BY APPLICANT
(Use Several Sheets If Necessary)

Applicant: Allen Comer et al.

(Use Several Sheets If Necessary)			Applicant: Allen Comer et al.	
7 CFR § 1.9	98(ъ))		Filing Date: 03/01/02	Group Art Unit:
		OTHER DOCUMENTS (Including Author, Title, D	Date, Relevant Pages, Place of Publication)	
Berger et al., Secreted placental Alkaline Phosphatas Gene 66:1-10 (1988)		Berger et al., Secreted placental Alkaline Phosphatase: A Pov Gene 66:1-10 (1988)	werful New Quantitative Indicator Of Gen	e Expression In Eukaryptic Cells,
Ĺ	2	Gibbs et al., Culture Of Reconstructed Epidermis In A Define And Improved Stratum Corneum, Arch Dermatol Res. 1997 S	ed Medium at 33°C Shows A Delayed Epid Sep;289(10):585-95. Erratum in: Arch Derr	dermal Maturation, Prlonged Lifes πatol Res 1998 Jan-Feb;290(1-2):2
	3.	Andraedis et al., Keratinocyte growth factor induces hyperprofileration and delays differtiation in a skin equivalent model system, FASEB 15:898-906 (2001)		
	4.	Auger et al., Multistep Production Of Bioengineered Skin Substitutes: SEquential Modulation Of Culture Conditions, In Vitro Cell Dev. BiolAnimal 36:96-103 (2000)		
	5.	Chilcott et al., Transepidermal Water Loss Does Not Correlate With Skin Barrier Function In Vivo, J. Investigative Dermatology 118:871-875 (2002)		
	6	Goretsky et al., Surface electrical capacitance as an index of epidermal barrier properties of composite skin substitutes and skin autografts Wound Repair and Regeneration 3:419-425 (1995)		
	7	Boyce et al., Surface Electrical Capacitance as a Noninvasive Index of Epidermal Barrier in Cultured Skin Substitutes in Athymic Mice, Soc. for Investigative Dermatology, 82-87 (1996)		
	8	Vicanova et al., Incorporation of linoleic acid by cultured human keratinocytes, Arch Dermatol Res. 291:405-412 (1999)		
	9	Swartzendruber et al., Molecular Models of the Intercellular Lipid Lamellae in Mammalian Stratum Comeum, Soc. for Investigative Dermatology,92:251257 (1989)		
	10	Punec et al., Lipid and ultrastructural scharacterization of reconstructed skin models, Int.I J. of Pharmaceutics, 203:211-225 (2000)		
	11	Gibbs et al., Temperature-sensitive regulation of epidermal morphogenesis and the expression of comified envelope precursors by EGF and TGFα, Cell Tissue Res., 292:107-114 (1998)		
	12	Uchida et al., Vitamin C Stimulates Sphingolipid Production and Markers of Barrier Formation in Submerged Human Keralinocyte Cultures, Soc. for Investigative Dermatology, 117:1307-1313 (2001)		
	13	Boyce et al., Vitamin C Reggulates Keratinocyte Viability, Epidermal Barrier, and Basement Membrane In Vitro, and Reduces Wound Concentration After Grafting of Cultured Skin Substitutes, Soc. for Investigative Dermatology, 118:565-572 (2002)		
-	14	Supp et al., Incubation of cultured skin substitutes in reduced mice, Wound Repair and Regeneration, 7:226-237 (1999)	humidity promotes comification in vitro a	nd stable engrastment in athymic
	15	Boyce et al., Lipid Supplemented Medium Induces Lamellar E Soc. for Investigative Dermatology, 101:180-184 (1993)	Bodies and Precursors of Barrier Lipids in	Cultured Analogues of Human SI
4n	16	Vicanova et al., Normalization of Epidermal Calcium Distribution Profile in Reconstructed Human Epidermis Is Related to Improvement of Terminal Differentiation and Straum Corneum Barrier Formation, Soc. for Investigative Dermatology, 111:97-106 (1998)		
				· · · · · · · · · · · · · · · · · · ·
aminer:	<u> </u>	in-Lin Chen	Du Carita in F) -1) t
arminel.		ial citation considered. Draw line through citation if not in conf	Date Considered:	U-03